

**DYNAMICS OF THE
YOUNG BINARY LMC CLUSTER NGC 1850**

PHILIPPE FISCHER¹, DOUGLAS L. WELCH¹

Department of Physics and Astronomy, McMaster University, Hamilton, Ontario L8S 4M1, Canada
Email: Fischer@Crocus.Physics.McMaster.CA

MARIO MATEO²

The Observatories of the Carnegie Institute of Washington
813 Santa Barbara Street, Pasadena, CA 91101

Address for proofs:

Philippe Fischer

Department of Physics and Astronomy

McMaster University

1280 Main Street West

Hamilton, Ontario

L8S 4M1 CANADA

Email: Fischer@Crocus.Physics.McMaster.CA

¹ Guest Investigator, Mount Wilson and Las Campanas Observatories, which are operated by the Carnegie Institution of Washington.

² Hubble Fellow.

Figures in postscript form are available via anonymous FTP as follows.
ftp 130.113.0.111 anonymous jemail address; cd pub get fig1850.ps quit

ABSTRACT

Isochrone fits to a BV color-magnitude diagram revealed that the primary cluster has an age of $\tau = 90 \pm 30$ Myr while the secondary member has $\tau = 6 \pm 5$ Myr. The reddening was found to be $E(B-V)=0.17$ mag. BV surface brightness profiles were constructed out to $R > 40$ pc, and single-component King-Michie (KM) models were applied. The total cluster luminosity varied from $L_B = 2.60 - 2.65 \times 10^6 L_{B\odot}$ and $L_V = 1.25 - 1.35 \times 10^6$ as the anisotropy radius varied from infinity to three times the scale radius with the isotropic models providing the best agreement with the data. Simple tests were made to check for tidal truncation in the profiles and we concluded that there was slight evidence favoring truncation. The bright background and binary nature of NGC 1850 render this conclusion somewhat uncertain.

Of the 52 stars with echelle spectra, a subset of 36 were used to study the cluster dynamics. The KM radial velocity distributions were fitted to these velocities yielding total cluster masses of $5.4 - 5.9 \pm 2.4 \times 10^4 M_\odot$ corresponding to $M/L_B = 0.02 \pm 0.01 M_\odot/L_{B\odot}$ or $M/L_V = 0.05 \pm 0.02 M_\odot/L_{V\odot}$. A rotational signal in the radial velocities has been detected at the 93% confidence level implying a rotation axis at a position angle of 100° . A variety of rotating models were fit to the velocity data assuming cluster ellipticities of $\epsilon = 0.1 - 0.3$. These models provided slightly better agreement with the radial velocity data than the KM models and had masses that were systematically lower by a few percent.

Values for the slope of the mass function were determined using the derived M/L , theoretical mass-luminosity relationships, and several forms for the IMF. The preferred value for the slope of a power-law IMF is a relatively shallow, $x = 0.29^{+0.3}_{-0.8}$ assuming the B-band M/L or $x = 0.71^{+0.2}_{-0.4}$ for the V-band.

1. INTRODUCTION

There is a large population of young, massive clusters in the LMC that have no counterparts in the Milky Way. These objects represent a unique opportunity to study the internal dynamics of resolved stellar systems in which the current ages are substantially less than the two-body relaxation timescales. The clusters can be studied kinematically

to determine masses and mass-to-light ratios (hereafter, M/L's) in order to constrain the initial mass function. Through the use of both imaging and spectroscopic data one can determine the value of various dynamical parameters and better understand formation and early cluster evolution.

To date there have been a number of dynamical studies of LMC clusters including integrated spectra for several old and intermediate age clusters (Elson and Freeman 1985; Dubath *et al.* 1990; Mateo *et al.* 1991; and Meylan *et al.* 1991), individual stellar radial velocity measurements of intermediate age clusters (Seitzer 1991, Fischer *et al.* 1992b) and individual velocity measurements for young clusters (Lupton *et al.* 1989, Mateo *et al.* 1991, Fischer *et al.* 1992a). Assuming stars can be resolved right to the cluster core, the individual velocity measurements are favored over the integrated spectra. With stellar radial velocities one can include radial information in the dynamical models, rotation can be detected and quantified, escaping stars can be detected, and, if multi-epoch observations are available, favorably aligned binary stars can be found.

NGC 1850 is a very bright young cluster in the Bar region of the LMC. It is located in a region rich with star clusters and star formation and is, in fact, embedded in the emission nebula Henize 103. Furthermore, the cluster appears to be in a binary or perhaps even a triple system, and there is a clear indication of tidal interactions. The time is ripe for both photometric and kinematic studies of this object. Only recently have complete stellar isochrones appeared for high mass stars enabling one to determine ages more reliably than in the past. The use of photon-counting detectors enables high-precision radial velocity measurements for stars with apparent magnitudes as faint as $V = 18$ mag.

In §2.1 the CCD imaging is described and, in §2.2, a color-magnitude diagram is constructed and analyzed in order to derive an accurate age estimate. In §2.3 surface density profiles are derived and in §3 King-Michie models are applied to this data. §4 contains a description of the radial velocity observations and reductions. §5 has the cluster mass estimates and §6 is a discussion of the cluster mass function. Finally §7 contains a

calculation of the relaxation timescales for the cluster.

2. CCD IMAGING

2.1 The Data

BV CCD frames of NGC 1850 were obtained at the Las Campanas Observatory (LCO) 1.0 m telescope on 1991 February 23. The TEK2 1024² chip was used (readout noise = 7 e⁻, gain = 2 e⁻/ADU, and angular scale = 0.61'' px⁻¹). The exposure times were 60 seconds for each filter.

There are many interesting objects in the immediate vicinity (i.e. within about 6 arcminutes) of NGC 1850 as can be seen from Figs. 1 and 2. Fig. 1 displays intensities around the median for the frame while Fig. 2 shows only the brighter stars. In Fig. 1 the cluster appears highly asymmetric with a large collection of stars slightly north of due east. This has been interpreted as both an interaction tail due to the presence of a binary companion (Bhatia and MacGillivray 1987) and as a cluster with a population distinct from the central cluster H88-159 (Bica *et al.* 1992, based on integrated photometry). Following the line joining the main cluster to this subcluster about 200 arcsec one finds a very low luminosity cluster (marked with a 1 on the figure). Approximately 100 arcsec south there is a diffuse object (2). Another, slightly higher luminosity cluster is located about 275 arcsec north of NGC 1850, and NGC 1855/54 (3), a bright young cluster, lies to the south, with just its edge visible in this frame (4). Another interesting feature is revealed in Fig. 2 directly to the West of NGC 1850 with a center-to-center separation of 30 arcsec (circled). This object has been previously designated by Bhatia & Hatzidimitriou (1988) as a distinct binary companion to NGC 1850 and according to Bica *et al.* is the probable source of ionizing radiation for the emission nebula Henize 103 in which these objects are embedded. Bhatia & Hatzidimitriou also concluded that the frequency with which apparent double clusters occurred in the LMC was much higher than one would expect from chance superpositions.

2.2 The Color-Magnitude Diagram

Stellar photometry was performed using DoPHOT (Mateo & Schechter 1989). As no standard stars were observed on this night the calibration was accomplished through the use of another set of NGC 1850 frames taken on 1990 November 5 using the same telescope, CCD and filters as above. On that night 12 BV observations of 9 E-region standards (Menzies *et al.* 1989) were observed covering an airmass range of 1.02 - 1.52 and a color range of $-0.043 \text{ mag} \leq B-V \leq 1.890 \text{ mag}$. The rms of the adopted solutions was approximately 0.02 mag. Aperture corrections yielded a zero-point with a similar accuracy and we conservatively estimate our absolute calibration to be accurate at better than 0.04 mag. Stars were then matched up between the two sets of frames to calibrate the present data. The reason for not using the older data for the current work is that the more recent frames had superior seeing which yielded improved stellar photometry particularly in the cluster's inner regions.

There have been at least three previous published photometric studies of NGC 1850. Robertson (1974) presented $(BV)_{pg}$ photographic photometry using an iris photometer and based his calibration on the $(BV)_{pe}$ photoelectric photometry of Tifft and Snell (1971). Alcaïno & Liller (1987) presented photographic BVRI photometry also using an iris photometer and based their calibration on a 15 star photoelectric sequence (Alcaïno & Liller 1982). Elson (1991) performed BV CCD photometry and based her calibration on four stars from Graham (1982), claiming a zero-point uncertainty of less than 0.01 mag. We have plotted the differences between our photometry and 44 stars from Robertson, 99 stars from Alcaïno & Liller, and 190 stars from Elson in Figs. 3 through 5, respectively. The mean zero-point differences, in the sense of us-them, are $\Delta V_{Robertson} = 0.29$, $\Delta B_{Robertson} = 0.36$, $\Delta V_{A\&L} = 0.06$, $\Delta B_{A\&L} = 0.03$, and $\Delta V_{Elson} = -0.14$, $\Delta B_{Elson} = -0.10$. We are fainter than Robertson and Alcaïno & Liller, brighter than Elson, redder than both Robertson and Elson and bluer than Alcaïno & Liller. There is a strong trend in both the B and V magnitude differences between ourselves and Robertson in the sense that the agreement

becomes systematically worse for the fainter objects. Since this trend is not significantly present for the other two comparisons we conclude that the Robertson data is incorrect.

Fig. 6 shows an apparent BV color-magnitude diagram (CMD) for the entire 10' square field. There is evidence for at least three distinct populations. Region 1 contains only very young stars still on the main sequence. Region 2 contains a mixture of the very young stars as well as a slightly older population of main sequence and evolved stars. Region three appears to consist of a significantly older population of evolved stars.

Fig. 7 contains a small subset of the stars found in Fig. 6. These are all the stars which were detected within 12 arcsec of the center of the binary companion located 30 arcsec to the west of the main cluster. In fact, there are many more stars in this region but unfortunately the inner region of the companion cluster was saturated on the 60s frames. Even with a shorter exposure we suspect that the extremely high degree of crowding in this region would render stellar photometry both difficult and unreliable. In any case, one can see that nearly all the detected stars are blue and they extend beyond the clearly delineated turnoff region obvious in Fig. 6 indicating a very young system.

We attempted to fit isochrones from Mermilliod *et al.* 1992 to the photometry data. These isochrones incorporate the new radiative opacities from Rogers & Iglesias (1992), convective overshooting and mass-loss (see Schaller *et al.* 1992 for a complete description of the stellar models). Isochrones were available for $z=0.0188$ (solar abundance), $z = 0.008$, and $z=0.001$. There are two published photometric metallicity determinations for NGC 1850: the first based on Washington photometry of two stars yielding $z = 0.009$ (Schommer & Geisler 1986), and the second based on Strömgren photometry of five stars yielding $z = 0.004$ (Grebel *et al.* 1992).

In Fig. 7 one can see the result of the isochrone fitting to the stars of the companion cluster. The inferred reddening was $E(B-V) = 0.17 \pm 0.03$ mag [$A_V = 3.1E(B-V)$ was assumed]. The presence of the young, barely-evolved, companion provides an excellent opportunity to accurately determine the reddening and our value agrees well with three

previous reddening determinations. Persson *et al.* (1983) find $E(B-V) = 0.15$ mag using the technique outlined in Cohen *et al.* (1981) involving both optical and IR broadband color indices. Alcaino and Liller (1987) obtained $E(B-V) = 0.18$ mag from a BVRI CMD study and Lee (1991) found $E(B-V) = 0.15 \pm 0.05$ for UBV photometry. The best age is $\tau = 6 \pm 5$ Myr. This changes by less than 1 Myr as one varies the distance modulus by ± 0.3 mag.

Fig. 8 is a CMD for the region within $1'$ of the center of the main cluster (it includes all the stars in Fig. 7). The apparent BV magnitudes for this inner region are presented in Table 1. Column 1, 6 and 11 are the stellar identifications, column 2, 7 and 12 are the X positions on the CCD (increases towards the south), column 3, 8 and 13 are the Y positions (increasing towards the east), columns 4, 9, and 14 are the apparent V magnitudes, and columns 5, 10, and 15 are the apparent B-V magnitudes.

The two younger isochrones in Fig. 8 are identical to those discussed above while the older set are for $\tau = 90$ Myr. We have adopted the reddening obtained above using the young stars. Our age estimate is therefore $\tau = 90 \pm 30$ Myr, where the error estimate takes into account the uncertainty in the zero-point and reddening and the difficulty in fitting the isochrones to this crowded data set. Varying the distance modulus by ± 0.3 affects the age estimate by approximately 10 Myr. Fig 9 shows population-synthesis Monte Carlo simulations based on the two different isochrones with an assumed photometric scatter of 0.05 mag and a mass function slope of 1.0 (see equation 15). One feature is immediately apparent in both cases, the lack of stars in the Hertzsprung gap compared to the observed data. Accepting the correctness of the models, there are at least three possible explanations for this effect: 1) The presence of young LMC field stars (and possibly a few stars from the young subcluster), 2) Binaries and close superpositions that were measured as single stars and 3) A possible time span for the initial star formation (this would need to be at least half the cluster age). We feel the first two effects are the likeliest and a close superposition of stars has been invoked by Welch & Stetson (1993) to explain the anomalously small

luminosity amplitude and high brightness of three cepheids in the crowded regions of the young LMC cluster NGC 1866. Another feature is that the evolution slows down around the red giant branch (RGB) and at the blue end of the horizontal branch (HB) causing stars to build up in these regions. This is observed and we conclude that the stars are more metal-poor than $z = 0.008$ based on a synthetic HB which does not extend far enough to the blue and a synthetic RGB which appears to be too red. The $z = 0.001$ isochrone appears to have too blue a RGB although it does provide a reasonable fit at the blue end of HB. Therefore, the metallicity lies between $z=0.008$ and $z=0.001$ but this conclusion is dependent on the adopted reddening and distance modulus.

Previous age estimates are $\tau = 40 \pm 10$ Myr (Hodge 1983, based on data from Tifft & Connolly 1973, and Robertson 1974), $\tau = 21 \pm 5$ Myr (Alcaino & Liller 1987) and $\tau = 40 \pm_{-10}^{+50}$ (Lee 1991). We have shown that the the Alcaino & Liller photometry and reddening estimate are both close to our own and therefore the different age estimates can be attributed to the different isochrones (they used the isochrones of Maeder & Mermilliod 1981) and the different age-determination techniques. Unfortunately, the photometry of Lee was not available for comparison.

Finally, we examined stars in the region of H88-159. Unfortunately, there are no stars in this region as bright as the turn-off for the main cluster. Therefore, it is impossible to derive an accurate age for this subcluster in order to determine if it is, in fact, a distinct population from the main cluster as indicated from the integrated photometry of Bica *et al.* (1992). An argument against it being distinct as opposed to being a tidal tail is that it appears to continue outward from the main cluster in a fairly straight line for about 2 arcmin (see Fig. 1).

2.3 Surface Photometry

Clearly, the presence of the above described sub-clustering coupled with the large number of LMC field stars located near NGC 1850 is going to complicate attempts to obtain reliable surface photometry. Thus, it is advantageous to remove the luminosity

contribution of any definite cluster non-members. DoPHOT was used to remove the stars in regions 1 and 3 (see Fig. 8) from both the B and V frames.

It is also favorable to determine the cluster ellipticity so that correspondingly elliptical apertures can then be used to perform the surface photometry. This is a difficult task for most globular clusters and particularly so for NGC 1850. In common with other clusters, NGC 1850 consists of resolved stars, but in this case the stars dominating the light are young supergiants. As well, there is the presence of the young binary companion. Fig. 10 shows a contour plot of the B and V star-subtracted, median filtered (filter radius of $9''$) images (solid lines). Superimposed on the contours are ellipses produced by the ELLIPSE task in the IRAF¹ STSDAS package which uses the ellipse fitting technique of Jedrzejewski (1987). Three things are immediately clear: 1) The ellipses do not provide a good model for the smoothed NGC 1850 light distribution, 2) the elliptical parameters change rapidly as a function of radius, and 3) there is rather poor agreement between the two different bandpasses. This leads us to believe that the ellipticities which we are measuring probably result from the presence of a number of bright stars which could not be adequately subtracted due to the extreme crowding. They are, therefore, not a good representation of the shape of the underlying mass distribution and hence, lacking anything better, we use circular contours to perform the surface photometry. We cannot, however, rule out the possibility that NGC 1850 is significantly elliptical. The use of circular apertures in such a case does not tend to result in systematic errors in the surface photometry, but does increase the photometric scatter (i.e., see Fischer *et al.* 1992b).

The cluster center was found using a multi-step procedure. The first step was to estimate the position of the center using the isophotes produced above. Surface photometry was then obtained utilizing this cluster center. A King model (see §3) was fit to the surface

¹ IRAF is distributed by the National Optical Astronomy Observatories, which is operated by the Association of Universities for Research in Astronomy, Inc., under contract to the National Science Foundation.

photometry and a new image was made with the fitted King parameters but with a smooth light distribution. We then cross correlated the artificial image with the original image to obtain small residual X and Y shifts. The final centers were within 2 arcsec of the original guesses.

Surface photometry was performed in a manner similar to Djorgovski (1988). The frames were broken up into a series of concentric circular annuli centered on the cluster. The annuli were further divided into eight azimuthal sectors. The average pixel brightness was determined for each sector in a given annulus and the *median* of the eight separate measurements was taken as the representative brightness at the area-weighted average projected radius of the annulus (i.e., the mean radius of all the pixels within the annulus which is approximately equal to the geometric mean). The standard error of the median of the eight sectors was adopted as the photometric uncertainty. Using the median as opposed to the more commonly adopted mean is essential in the case of NGC 1850 as it reduces the contamination due to the bright supergiants and the binary companion.

A background level (a combination of sky light and Galactic foreground and remaining LMC field stars) was estimated from regions at large projected distances from the cluster. We found that the surface brightness profiles tended to level out beyond 45 pc (we have adopted a distance to the cluster of 50 kpc) for both the B and V frames. By “levelling out” we don’t necessarily mean that the cluster light does not extend beyond this point but simply that fluctuations in the background dominate to such an extent that it is no longer possible to observe the profile declining in intensity. Therefore, it was this region with a projected radius of $45 \leq R \text{ (pc)} \leq 100$ that was used for the background determinations. The reddening-corrected background values for the B and V frames were, respectively, $1370 \pm 27 L_{B\odot} \text{ pc}^{-2}$ and $682 \pm 14 L_{V\odot} \text{ pc}^{-2}$.

The background-subtracted surface photometry data is presented in Table 2 [assuming a cluster distance of 50 kpc, $M_{V\odot} = 4.83$ and $(B-V)_{\odot} = 0.65$, Mihalas and Binney 1981, p. 60]. Columns 1 and 3 are the projected area-weighted radii, and columns 2 and 4 are

the B and V luminosity densities, respectively.

Fig. 11 is a plot of the B and V surface brightness profiles. Also shown are typical stellar profiles which have a FWHM less than 15% of the cluster core radius (see §3); hence seeing will have a negligible effect on measurements of this quantity (Mihalas and Binney 1981, p. 315).

3. KING-MICHIE MODELS FOR THE SURFACE PHOTOMETRY

We fit projected single-component King-Michie (KM) models (King 1966 and Michie 1963) to the surface photometry data. These models have an energy (E) angular momentum (J) per unit mass distribution function given by

$$f(E = -0.5v^2 + W, J) \propto e^{-[J/(2v_s r_a)]^2} (e^{-0.5v^2 + W} - 1), \quad (1)$$

where v_s is the scale velocity, r_a is the anisotropy radius (both described below), and W is the reduced gravitational potential. The shape of the density distribution, $\rho_K(r)$, is determined by solving Poisson's equation and is dependent on two parameters; the central potential W_o , and r_a , beyond which stellar orbits become increasingly radial. Scaling is applied in both the radial (r_s) and luminosity (ρ_{K_o}) dimensions to give the best fit. For a complete description of the models (albeit for the more complex multi-component case) the reader is directed to Gunn and Griffin (1979). Model density profiles with r_a values ranging from $3 r_s$ to infinity (i.e., an isotropic distribution function) were generated, and projected on to the observational plane. These were binned identically to the observed data and fit using a maximum likelihood technique.

Tables 3 and 4 show, for each r_a , the best *fitted* KM parameters for both bandpasses. Column 1 contains r_a , column 2 is the reduced central potential, column 3 is r_s and column 4 is $c = r_s/r_t$ (r_t is the tidal radius). Column 5 is the reduced chi-squared ($\chi_\nu^2 = \chi^2/\nu$, where $\nu = 18$ is the number of degrees of freedom) for the fit. Column 6 is the probability of obtaining a value greater than χ_ν^2 for a model with the given parameters and the uncertainties listed in Table 2. These were derived from 1000 simulations of the surface

photometry data per model. Each simulation used a surface profile generated from the best fit model with errors, drawn from the uncertainties shown in Table 2. The same fitting procedure originally applied to the real data was utilized and, in this way, we found the uncertainties in each fitted parameter as well as the distribution of χ^2_ν . The remaining columns of Tables 3 and 4 will be discussed in §5.1. The fitted KM model parameters were consistent for the two bandpasses.

Tables 5 and 6 have a summary of the *derived* KM parameters for both bandpasses corresponding to the models specified in Tables 3 and 4. Column 1 is the anisotropy radius while column 2 is the central luminosity density, $\rho_{K\odot}$. Column 3 is the cluster luminosity and columns 4 and 5 will be discussed in §5.1.

The solid lines in Fig. 11 show the best fit models which are those possessing isotropic orbits. The quality of the fits deteriorates as r_a decreases. Interestingly, this is quite different from what was seen in the case of the slightly older LMC cluster NGC 1866 which favored highly anisotropic (i.e. $r_a = 3 r_s$) KM models and appeared to have a halo of unbound stars (Fischer *et al.* 1992a). Also shown as dotted lines in the same figure are the $r_a = 3 r_s$ models. It is unclear whether or not the data exhibit a tidal cut-off. In fact, using a profile that extends out to about 20 pc, Elson (1991) finds no evidence for truncation. Our profile extends to about twice that radius and there is, at best, a small indication for truncation. To test the hypothesis that there is significant truncation we have fit models of the form:

$$\mu(R) = \mu_\odot [1 + (R/a)^2]^{-\gamma/2}, \quad (2)$$

to the surface brightness profile using a non-linear, weighted, least squares technique. The best-fit parameters for these untruncated power-law models are: $\mu_{B\odot} = 26800. \pm 2000. \odot \text{ pc}^{-2}$, $a_B = 2.8 \pm 0.3 \text{ pc}$, $\gamma_B = 2.26 \pm 15$, $\mu_{V\odot} = 12000. \pm 900. \odot \text{ pc}^{-2}$, $a_V = 3.1 \pm 0.4 \text{ pc}$, $\gamma_V = 2.34 \pm 0.15$. The χ^2_ν are 0.89 [P(> χ^2_ν) 0.59] and 1.06 [P(> χ^2_ν) 0.39] for the B and V profiles, respectively, which are higher than all but the $r_a = 3 r_s$ KM models.

We conclude that the best truncated models provide marginally better agreement with the surface photometry data.

4. RADIAL VELOCITIES

4.1) Observations and Reductions

Spectra of 52 supergiants in the region surrounding NGC 1850 were obtained during two runs (1991 February 14-20 and 1991 December 14-17) using the photon-counting echelle spectrograph on the 2.5m Dupont reflector, designed and built by Steve Shectman. Eight of the stars have repeat measurements.

The observation and reduction procedures for a previous run at LCO have been discussed extensively in Welch *et al.* (1991) and remain largely unchanged for this data. Briefly, the observing procedure consisted of exposures with integration times of 200 - 500s and Th-Ar arcs approximately every 45 minutes. A representative LCO spectrum is shown in Fig. 2 of Côté *et al.* (1991). The reduction utilizes the IRAF ECHELLE and RV packages (Tody 1986) to obtain both velocities and velocity uncertainties. The velocity zero-point is tied to the IAU velocity standard 33 Sex as described in Fischer *et al.* 1992a and is believed to be accurate to better than 2 km s^{-1} .

Along with the program stars, relatively high S/N spectra were obtained on each night for both a nearby bright star used as a local velocity standard (LVS) for differential measurements and the radial velocity standard HD 23214 to examine the possibility of velocity zero-point drifts and to test the accuracy of the velocity uncertainties returned by RVXCOR. These velocities are shown in Table 7 for the eight separate nights over the two observing runs in which NGC 1850 stars were observed. Column 1 is the Heliocentric Julian Date minus 2448000, column 2 is the radial velocity, column 3 is the mean velocity for the given observing run and column 4 is the reduced chi squared for the appropriate observing run. Within the RVXCOR produced uncertainties, there is no significant zero-point drift during the individual runs. There is, however, a discrepancy between the mean velocities for the LVS for the two separate observing runs at the 4.6σ level implying either

that the radial velocity zero-point has shifted or that this star has a variable velocity. The shift is not seen in HD 23214 implying the latter hypothesis is probably correct. Olszewski *et al.* (1991) have obtained 9 lower resolution radial velocity measurements for HD 23214 during two observing runs, 1987 January 8-12 and 1987 December 27-31, deriving a mean velocity of $-4.3 \pm 1.8 \text{ km s}^{-1}$. They also present a single CORAVEL velocity determination of $-4.7 \pm 0.3 \text{ km s}^{-1}$ for HJD = 2446862.51. This latter deviates by 5.6σ from our value of $-2.8 \pm 0.2 \text{ km s}^{-1}$, which, as mentioned above, is based upon the IAU radial velocity standard 33 Sex. Another radial velocity standard, HD 196983, has been measured at two epochs, HJD - 2448000 = 606.5225 and 608.5179 yielding $v_x = -9.6 \pm 0.5 \text{ km s}^{-1}$ and $-8.7 \pm 0.6 \text{ km s}^{-1}$, respectively, for a mean of $-9.2 \pm 0.4 \text{ km s}^{-1}$. This agrees very well with the CORAVEL value, based on 10 separate measurements spanning 715 days starting in September 1981, of $-9.3 \pm 0.1 \text{ km s}^{-1}$ (Maurice *et al.* 1984). To conclude, we feel that our velocity zero-point is uncertain at about the 2.0 km s^{-1} but that there is no significant zero-point drift during the two runs. (This latter possibility will be examined in more detail presently.)

The radial velocity data for the cluster stars are presented in Table 8. Column 1 contains the stellar identifications (these do not correspond to the ID's in Table 1), column 2 has the projected radii, column 3 the equinox J2000.0 position angles, column 4 contains the radial velocities and column 5 contains the mean velocities for stars with repeated measurements. Column 6 is the Heliocentric Julian Date - 2448000 for the velocity measurements. Columns 7 and 8 are V and B-V for the stars. The photometry in the innermost regions is relatively uncertain due to the high degree of crowding. Fig. 12 is a finder chart for all but the star farthest from the cluster center.

Of the eight stars with repeated velocity measurements, six had individual χ^2 of less than 2.85 for a total $\chi^2 = 6.44$ for 5 degrees of freedom. Of the other two, one had $\chi^2 = 7.9$ (star RV 49) and the other had $\chi^2 = 217$ (star RV 19). If the velocity shift indicated by the LVS is applied these χ^2 values all decrease; we get a total $\chi^2 = 4.33$ for the six stars

and 3.40 and 217.00 for the other two stars. This is an argument in favor of adoption of the velocity shift.

From their colors, the two high- χ^2 stars are evolved and their velocities indicate that they are in the LMC. Star RV49 appears to be an LMC field star, perhaps a binary. Star RV19, as indicated by its radial velocities, is clearly a member of the LMC. Further, its luminosity ($V = 13.54$) indicates that it is a supergiant. Two possibilities are that the star is a binary or a Cepheid. The two velocity measurements differ by over 30 km s^{-1} which argues against the former hypothesis as it would require a very favorable inclination coupled with fortuitous observations. The stars color ($B - V = 0.65$) makes it an excellent Cepheid candidate, and it is likely to be a cluster member based on its proximity to the cluster center ($R = 3.8 \text{ pc}$). Neither of these two stars will be used in the cluster mass determinations.

In Fig. 13 we present plots of radial velocity and radial velocity uncertainty vs. stellar $B-V$. There are clearly trends towards both a larger apparent velocity dispersion and larger uncertainties for the bluer stars. The correlation between velocity uncertainty and apparent velocity dispersion is, of course, expected. The correlation between uncertainty and color has two causes: 1) the lack of lines in the bluer stars, and 2) spectral mismatch with the template star (a K giant). For the mass determinations of the next section we decided not to employ any stars with uncertainties greater than 2.7 km s^{-1} (the solid lines in Fig. 13), eliminating 9 stars. We have also eliminated the three stars having $v_x > 270 \text{ km s}^{-1}$ as being probable non-members (or variables) and star RV 18, which was classified as a variable by Robertson (1974) and from its color is a probable cepheid, leaving a total of 38 stars.

5. VELOCITY DISTRIBUTIONS AND MASS DETERMINATIONS

5.1 King-Michie Models

The mass of a multi-mass KM model is given by

$$M = \frac{9r_s v_s^2}{4\pi G} \int \frac{\rho}{\rho_0} r^2 dr \quad (3)$$

Illingworth (1976), where r_s is given in Tables 3 and 4, and v_s is the scale velocity. The run of $\sigma_r^2(r)$ and $\sigma_t^2(r)$ are determined from

$$\sigma_{r,t}^2(r) = \frac{\int_{\sigma \leq W(r)} f(\sigma, W) \sigma_k^2 d^3 \vec{\sigma}}{\int_{\sigma \leq W(r)} f(\sigma, W) d^3 \vec{\sigma}}, \quad (4)$$

where W is the reduced potential ($W = 0$ at the tidal radius) and $\sigma_k = \sigma \cos \theta$ or $\sigma \sin \theta$ for σ_r or σ_t , respectively. Comparisons were made between the observed velocities and the model velocity dispersion projected along the line of sight,

$$\sigma_p^2(R) = \frac{2}{\mu(R)} \int_R^\infty \frac{\rho_K(r) [(r^2 - R^2) \sigma_r^2(r) + R^2 \sigma_t^2(r)] dr}{r(r^2 - R^2)^{1/2}}, \quad (5)$$

(Binney and Tremaine 1987, p. 208), yielding the optimal value for v_s . The comparison was accomplished using the maximum likelihood technique outlined in Gunn and Griffin (1979). Simply put, the probability density function for v_{xi} , an observed stellar velocity, is a Gaussian with standard deviation equal to the model dispersion plus the velocity uncertainty added in quadrature:

$$P_i \sim \frac{1}{\sqrt{v_{err\ i}^2 + v_{p\ i}^2}} e^{-(v_{x\ i} - v_{ave})^2 / 2(v_s^2 v_{p\ i}^2 + v_{err\ i}^2)}. \quad (6)$$

One minimizes this function with respect to v_s and v_{ave} resulting in two equations which can be solved simultaneously for the most probable values of the two parameters.

A serious problem in mass determinations is contamination from binary and non-member stars, both of which tend to increase the mass estimate. This is particularly a problem when one has a relatively small sample of stars and a small velocity dispersion as is the case for NGC 1850. A single interloper in the data sample can easily result in a 50% or greater overestimate in the mass. We attempted to deal with this problem in the

following manner. First, using the entire data set, the optimal v_s and v_{ave} are determined using equation 6. For every star the parameter

$$\delta_i = \sqrt{\frac{(v_{ri} - v_{ave})^2}{v_s^2 \sigma_p^2(R_i) + v_{err\ i}^2}} \quad (7)$$

is tabulated. The star with the largest δ_i is removed and the procedure is repeated until all the stars have been removed. Then, using the KM models, we performed Monte Carlo simulations of the radial velocity data. We started with the known projected radii (R_i) of the program stars. The true radius is in the range $R \leq r \leq r_{max}$, where r_{max} can extend to infinity for an unbound distribution. If x is the displacement from the mean cluster position along the line-of-sight such that $r = \sqrt{R^2 + x^2}$ then the probability that the star is at x is

$$p(x) \sim \rho_K(\sqrt{R^2 + x^2}). \quad (8)$$

A three-dimensional position along with corresponding model-dependent radial and tangential velocities were drawn at random from their respective probability distributions. The velocity component along the line-of-sight was then determined, and an error term, drawn from a Gaussian distribution with standard deviation equal to the velocity error, as tabulated in Table 8, was added. This process was repeated, producing 10000 sets of data, each with a given mass and r_a and the same projected positions and velocity measurement errors as the original data set. Finally the maximum likelihood technique was applied to each of the artificial data sets and the maximum δ_i were recorded. The v_s for the first three iterations (using the unshifted radial velocity data) are shown in column 2 of Table 9. Columns 3 and 4 are the maximum δ_i and the percentage of simulations with δ_{max} exceeding this value, respectively, and column 5 is the star possessing δ_{max} . A value of zero in the fourth column indicates less than 0.001. We feel that there is a fairly high probability that the first two stars are either variables or non-members and hence they will not be used for the mass determinations. It is worth remembering, however, that their removal results in a mass reduction of about 80% demonstrating the extreme sensitivity

of the mass determinations to interlopers. We repeated the procedure with data that was corrected according to the LVS (see §4.1) and found similar results but obtained a v_s that was about 2% larger. The fact that it causes an increase in the velocity dispersion argues against its use, but in any case it does not have significant repercussions on the mass estimates and hence was not adopted.

Fig. 14 shows mean radial velocity vs. projected radius (upper panel) and versus position angle (lower panel) for the 36 remaining stars. The solid lines are the mean velocity, $\bar{v} = 251.4 \pm 2.0 \text{ km s}^{-1}$.

The values of v_s obtained from this reduced data set are displayed in column 7 of Tables 3 and 4. The corresponding masses and M/L ratios are in columns 4 and 5 of Tables 5 and 6. One can see that the total cluster mass and M/L are fairly insensitive to assumptions about r_a , and the best values are $M = 5.7 \pm 2.3 \times 10^4 M_\odot$ and $M/L_B = 0.02 \pm 0.01 M_\odot/L_{B\odot}$ or $M/L_V = 0.05 \pm 0.02 M_\odot/L_{V\odot}$.

The above-described Monte-Carlo orbit simulations were used to determine the uncertainties implicit in the maximum likelihood technique and to search for any possible systematic effects. We found that this method tended to underestimate v_s by about 3% and hence the mass by 6%. The values in the relevant tables have been corrected for this effect and the uncertainties shown were derived from the simulations.

A goodness-of-fit statistic

$$\chi^2 = \sum \frac{(v_{x\ i} - v_{ave})^2}{(v_s^2 v_{p\ i}^2 + v_{err\ i}^2)} \quad (9)$$

was generated for each value of r_a and is shown in column 8 of Tables 3 and 4 (34 degrees of freedom). The distribution of this statistic can be extracted from the Monte Carlo simulations. Column 9 shows the probability of exceeding the observed χ^2 assuming that the cluster velocities are specified by the model parameters indicated and have the uncertainties tabulated in Table 8. The isotropic models yielded the best agreement with the data (although, only marginally so) consistent with the findings from the surface photometry.

Because of the similarity between the velocity dispersion and the velocity uncertainty, the mass estimates are quite highly dependent on the accuracy of the uncertainty estimates. We have already discussed the analysis of the stars with repeated measurements and concluded that the uncertainties seem to be reasonable with reduced χ^2 near unity. However, it is worth quantifying the effects of the uncertainties. Assuming that the uncertainties are actually zero causes an increase in the estimates of the v_s 's by approximately 25%, or a roughly 50% increase in cluster mass. Increasing the uncertainties by 50% lowers the v_s 's by 50% and the mass by 75%.

5.2 Rotation

A careful examination of the radial velocities vs. position angle (Fig. 14) reveals evidence for a sinusoidal variation which may be indicative of cluster rotation. In order to test this hypothesis we measured the difference in median velocities on either side of an imaginary axis which is stepped around the cluster center at 1° intervals. If one is viewing an edge-on rotating system then a sinusoidal variation is expected with the velocity difference maximized when the axis corresponds to the rotation axis. This effect will degrade as the system deviates from edge-on. The bottom left panel of Fig 15 shows a plot of the velocity differences versus the position angle of the axis. The best-fit sine curve (using unweighted least-squares) has an amplitude of $A = 2.1 \text{ km s}^{-1}$ and the implied projected rotation axis is $100^\circ \pm 40^\circ$ ($\chi^2 = 25.12$ for 178 degrees of freedom). To test the significance, we constructed 1000 non-rotating models as described above and performed the same test. Only 7% of these models had amplitudes exceeding $A = 2.1$ and hence we feel fairly confident that NGC 1850 is rotating. Rotation has previously been detected at the 97% confidence level for a sample of 69 stars in the young LMC cluster NGC 1866 (Fischer *et al.* 1992a) which had $A = 1.8 \text{ km s}^{-1}$ and $v_s = 3.1 \text{ km s}^{-1}$. In that case, incorporation of rotation resulted in only a very small downward change in the mass estimate. However, in the case of NGC 1850, the two parameters have very similar values meaning rotation may be dynamically more significant than it was in NGC 1866.

Unfortunately, as explained in §2.3 it is impossible to extract a meaningful ellipticity from the NGC 1850 light distribution. This information would have enabled us to construct rotating oblate cluster models to compare with the radial velocities. We therefore adopted an alternative approach of assuming different ellipticities values for the cluster, constructing models for the rotation, and comparing them to the data.

For an axisymmetric system, the relevant Jeans' equations (velocity moments of the collisionless Boltzmann equation) in cylindrical coordinates are:

$$\frac{\partial(\rho\sigma_R^2)}{\partial R} + \frac{\partial(\rho\sigma_{Rz})}{\partial z} + \rho \left(\frac{\sigma_R^2 - \sigma_\phi^2 - v_\phi^2}{R} \right) + \rho \frac{\partial\Phi}{\partial R} = 0, \quad (10)$$

and

$$\frac{\partial(\rho\sigma_{Rz})}{\partial R} + \frac{\partial(\rho\sigma_z^2)}{\partial z} + \frac{\rho\sigma_{Rz}}{R} + \rho \frac{\partial\Phi}{\partial z} = 0, \quad (11)$$

where (R, ϕ, z) are the cylindrical coordinate axes, the $(\sigma_R, \sigma_\phi, \sigma_z)$ are the corresponding velocity dispersions, and v_ϕ is the rotation velocity. Φ is the gravitational potential.

Both the rotating and non-rotating models which were used have velocity ellipsoids aligned with the cylindrical coordinate axes (i.e. $\sigma_{Rz} = 0$) and, as well, both have

$$\sigma_\phi = \frac{\sigma_R}{\sqrt{1 + (R/R_a)^2}}, \quad (12)$$

where R_a can be varied up to ∞ . The rotating models also have the condition

$$\sigma_R = \sigma_z, \quad (13)$$

implying

$$v_\phi^2 = R \frac{\partial\Phi}{\partial R} + \frac{R}{\rho} \frac{\partial}{\partial R} \int_z^\infty \rho \frac{\partial\Phi}{\partial z} dz + \frac{1}{\rho} \left[1 - \frac{1}{\sqrt{1 + (R/R_a)^2}} \right] \int_z^\infty \rho \frac{\partial\Phi}{\partial z} dz. \quad (14)$$

The models are constructed by assuming that the mass distribution is equivalent to the deprojected light distribution (constant M/L). Equations 11 and 14 can then be solved directly to obtain σ_z , and v_ϕ . Once v_ϕ is known it can be substituted into equation 10 which, in turn, can be solved for σ_R and σ_ϕ . This is outlined in Binney and Tremaine

(1987, p. 209) for the $\sigma_R = \sigma_\phi = \sigma_z$ case and a $\epsilon = 0.3$ model is shown in Fig. 6 of Fischer *et al.* (1992b).

Once the models have been generated, it is simply a matter of projecting and then scaling them using a similar maximum likelihood method to that employed for the KM models. For our purposes we have constructed models spanning $\epsilon = 0.1 - 0.3$ for $R_a = \infty$. All of the models assume a cluster inclination of 90° . Table 10 displays the results of the model fitting. Column 1 is the ellipticity, column 2 the cluster mass, columns 3 and 4 the B and V M/L's, and columns 5 and 6 are ζ^2 and $P(> \zeta^2)$ as above. These models all have $P(> \zeta^2)$ greater than or equal to the best non-rotating KM models despite having larger ζ^2 values. However, as can be seen from Fig. 7 of Fischer *et al.* (1992b) the rotating models have a wider ζ^2 distribution. The rotating models have masses (and M/L's) which are marginally lower than the non-rotating models. Fig. 15 exhibits plots of the velocity differences vs. axis position angle for the original and rotation-subtracted data. Column 7 of Table 10 is the amplitude of the best-fit sine curves for each case. One can see that the amplitude decreases for the rotation subtracted data with increasing ϵ . It does not, however, decrease smoothly to zero and one can see that for the $\epsilon = 0.3$ case there is a residual trend in the data which does not disappear for higher ellipticities. This probably results from some type of model mismatch which may be a result of the model itself or might be due to a poorly determined rotation axis position angle or an inclination of less than 90° . We changed the rotation axis position angle by plus and minus 20° but found that the residual velocities did not improve. Another possibility is that the characteristic rotational signature arises from the interaction between the two binary cluster components. The resulting stellar motions might in such a case have a similar appearance to rotation.

6. CONSTRAINTS ON THE MASS FUNCTION

The cluster M/L estimate can be used to constrain the slope of the initial mass function (IMF). In this study we used an IMF of the form

$$\phi(m) = m^{-(x+1)} dm \quad m \geq m_d, \quad (15)$$

$$\phi(m) = m \, dm \quad m < m_d, \quad (16),$$

which gives a drop-off at the faint end similar to what is seen in the solar neighborhood (Miller & Scalo 1979).

The theoretical cluster M/L is given by

$$\frac{M}{L} = \frac{\int_{m_l}^{m_u} m \phi(m) dm}{\int_{m_l}^{m_u} l(m) \phi(m) dm}, \quad (17)$$

where $l(m)$ is the luminosity of a star of mass m given by a theoretical mass-luminosity relationship for main-sequence and evolved stars. We used the mass-luminosity relationship of Mermilliod (1992), supplemented with the Bergbusch & Vandenberg (1992) values in the range $0.15 \leq m \, (M_\odot) \leq 0.9$, with the cluster parameters determined in §2.2. The initial mass of a star now at the end of the asymptotic giant branch is approximately $5.4 \, M_\odot$. We have adopted the treatment of Pryor *et al.* 1986 to deal with remnants: stars with initial masses of $5.4 - 8.0 \, M_\odot$ become white dwarfs with masses of $1.2 \, M_\odot$ and stars with initial masses greater than $8 \, M_\odot$ are assumed to be ejected from the cluster as is consistent with the high velocities seen for pulsars in the disk (Gunn and Griffin 1979). Therefore, the choice of m_u is not important except for estimating the amount of mass lost from the cluster due to stellar evolution.

Table 11 shows the derived values of x (column 3 and 5) for 3 different m_l (column 1), and 5 different m_d (column 2). Columns 4 and 6 contain the implied mean stellar mass for the given model. The values of x correspond to the B and V M/L's indicated at the top of the table. Although the values corresponding to the B and V M/L's are consistent within the uncertainty, the B-band x 's are systematically lower. It appears that for a given set of assumptions the V-band M/L constrains the mass function slope somewhat more tightly than the B-band. Furthermore, the V-band is also less sensitive to the assumed cluster age although one finds that the both the slopes steepen for a younger assumed age and flatten for older (this has not been accounted for in the uncertainty). The B-band has one advantage in that it appears to be less sensitive to the low mass cut-off. Both sets of slopes

are substantially shallower than was seen for the young LMC cluster NGC 1866 (Fischer *et al.* 1992a), which had an average slope (i.e. as determined without employing a drop-off in the mass function) of between $x = 1.35 \pm 0.1$ and $x = 1.82 \pm 0.1$ (depending on adopted age) for $m_l = 0.1$.

7. Evolutionary Timescales, Mass Segregation and Equipartition of Energy

The stars with measured velocities are all supergiants, with relatively large masses. If equipartition of energy has occurred these stars, being the most massive, would have a velocity dispersion below the mean value for all mass classes. Therefore, in order to justify the use of single-mass models we must demonstrate that the cluster is sufficiently young such that equipartition and mass segregation have not had enough time to become significant effects. Two important timescales are the central relaxation time

$$t_{ro} = (1.55 \times 10^7 \text{yr}) \left(\frac{r_s}{\text{pc}} \right)^2 \left(\frac{v_s}{\text{km s}^{-1}} \right) \left(\frac{M_\odot}{\langle m \rangle} \right) [\log(0.5M/\langle m \rangle)]^{-1} = 0.4 - 2.2 \times 10^8 \text{yr} \quad (18)$$

(Lightman and Shapiro 1978) and the half mass relaxation time

$$t_{rh} = (8.92 \times 10^8 \text{yr}) \left(\frac{M}{10^6 M_\odot} \right)^{1/2} \left(\frac{r_h}{\text{pc}} \right)^{3/2} \left(\frac{M_\odot}{\langle m \rangle} \right) [\log(0.4M/\langle m \rangle)]^{-1} = 1.5 - 5.3 \times 10^9 \text{yr} \quad (19)$$

(Spitzer and Hart 1971). The parameters in these equations have all been discussed previously except for the half mass radius, $r_h \simeq 11$ pc. Aside from the inner core the cluster is in a dynamically unevolved state and, therefore, we do not expect substantial energy transfer to have occurred. We conclude that no large systematic errors are being introduced into the velocity dispersions.

Another possible problem is primordial mass segregation. This would occur if star formation in the dense core region favored a different ratio of high-to-low mass stars than at larger radii. While this should not effect the velocity dispersion greatly, it would mean that the assumption of a uniform M/L is incorrect. Consequently the derived luminosity density profiles would not be an adequate representation of the mass density profile. One

should be able to construct luminosity functions at different projected cluster radii and hence search for a gradient in mass function. In practice, owing to the very crowded nature of the inner cluster regions and the high surface density of non-member stars present on the frame this would be difficult to accomplish in a convincing manner and we have not attempted it.

VII. CONCLUSIONS

In this paper we have examined the age and internal dynamics of the young binary LMC cluster NGC 1850 using BV CCD images and echelle spectra of 52 supergiants.

- 1) A BV CMD was constructed for the field surrounding the cluster and was found to contain 3 distinct populations of stars. The first was a very young population of age $\tau = 6 \pm 5$ Myr belonging primarily to the smaller member of the binary system located $30''$ west of the larger member. This young population allowed for an accurate reddening estimate of $E(B-V) = 0.17 \pm 0.03$ mag. The second, slightly older population, belongs primarily to the the larger cluster and has an age of $\tau = 90 \pm 30$ Myr. The third population was older and mainly comprised of LMC field stars.
- 2) Attempts were made to determine ellipticity parameters for the cluster using star-subtracted, median filtered BV images. This was greatly complicated by the presence of extremely bright young resolved stars and the binary nature of the cluster and no meaningful shape parameters were derived.
- 3) BV luminosity profiles were constructed out to projected radii of $R > 40$ pc. Single component King-Michie (KM) models were applied to this data in order to determine the most favorable value for the anisotropy radius and the total cluster luminosity. The luminosity varied from $L_B = 2.60 - 2.65 \pm 0.2 \times 10^6 L_{B\odot}$ and $L_V = 1.25 - 1.35 \pm 0.1 \times 10^6 L_{V\odot}$ as r_a went from to infinity to $3r_s$. The fitted and derived KM parameters for both bandpasses were consistent and the isotropic models provided the best agreement with the data.
- 4) To test for the presence of a tidal cut-off in the luminosity profile, a power-law

model without truncation was applied to the data. This model was found to provide marginally worse fits than the KM models giving some indication that the cluster density distribution is truncated.

- 5) Of the 52 stars with echelle spectra, a subset of 36 were used to study the cluster dynamics. The KM radial velocity distributions were fitted to these velocities yielding scale velocities of $v_s = 2.2 - 2.3 \pm 0.5 \text{ km s}^{-1}$ for the r_a range employed. The total cluster mass was $5.4 - 5.9 \pm 2.4 \times 10^4 \text{ M}_\odot$ corresponding to $M/L_B = 0.02 \pm 0.01 \text{ M}_\odot/L_{B\odot}$ or $M/L_V = 0.05 \pm 0.02 \text{ M}_\odot/L_{V\odot}$. The mean cluster velocity is $\bar{v} = 251.4 \pm 2.0 \text{ km s}^{-1}$.
- 6) A rotational signal in the radial velocities has been detected at the 93% confidence level implying a rotation axis at a position angle of $100^\circ \pm 40^\circ$. A variety of rotating models were fit to the velocity data assuming $\epsilon = 0.1 - 0.3$. These models provided slightly better agreement with the radial velocity data than the KM models and had masses that were systematically lower by a few percent.
- 7) Values for the slope of the mass function were determined using the derived M/L, theoretical mass-luminosity relationships, and several forms for the IMF. The preferred value for the slope of a power-law IMF is a relatively shallow, $x = 0.29^{+0.3}_{-0.8}$ assuming the B-band M/L or $x = 0.71^{+0.2}_{-0.4}$ for the V-band.
- 8) The current cluster age is similar to its central relaxation time but about 2 orders of magnitude less than its half-mass relaxation time. Therefore, aside from in the inner core the cluster is in a dynamically unevolved state and we expect that equipartition has not yet occurred in any substantial way.

ACKNOWLEDGEMENTS

P.F. would like to acknowledge the Natural Sciences and Engineering Research Council (NSERC) for a post-graduate fellowship. This work was undertaken while D.L.W. was a NSERC University Research Fellow. Partial support for this work was provided by NASA through grant # HF-1007.01-90A awarded by the Space Telescope Science Institute which

is operated by the Association of Universities for Research in Astronomy, Inc., for NASA under contract NAS5-26555. We would like to thank Dr. Maeder, Dr. Meynet, Dr Schaller, and Dr. Schaerer for a copy of their new isochrones and Dr. P. Bergbusch for a copy of his isochrones, both, prior to publication.

Reprints of this paper are available through mail or anonymous FTP. Contact Fischer@crocus.physics.mcmaster.ca for information.

FIGURE CAPTIONS

Fig. 1– A B band CCD image of NGC 1850 displayed to highlight fainter features. The image is approximately $10'$ square with north at the top and east to the left. See text for an explanation of the numbers and circles.

Fig. 2– The same as Fig. 1 but displayed to show only the bright stars.

Fig. 3– Comparison between photometry from this work and Robertson (1974). The zero-point differences can be found in the text.

Fig. 4– The same as figure 3 but for the photometry from Alcaïno & Liller (1987).

Fig. 5– The same as figure 3 but for the photometry from (Elson 1991)

Fig. 6– BV color-magnitude diagram for the entire $10'$ field surrounding NGC 1850. The numbers correspond to three distinct populations (see text).

Fig. 7– BV color-magnitude diagram for the region within $12''$ of center of the companion cluster (see Fig. 2). The solid line corresponds to the isochrone with $z = 0.008$ while the broken line has $z = 0.001$. Both sets of isochrones have ages of $\tau = 6$ Myr, as well as $(m-M)_0 = 18.5$ and $E(B-V) = 0.17$ mag.

Fig. 8– BV color-magnitude diagram for the region within $1'$ of the center of the main cluster. The solid lines correspond to the isochrones with $z = 0.008$ while the broken lines have $z = 0.001$. Both sets of isochrones have ages of $\tau = 6$ Myr and $\tau = 90$ Myr, as well as $(m-M)_0 = 18.5$ and $E(B-V) = 0.17$ mag.

Fig. 9– Population synthesis Monte Carlo experiments for the $z = 0.008$ and $z = 0.001$ isochrones described in the previous figure. A photometric uncertainty of 0.05 mag has been added.

Fig. 10– Contour plot of star-subtracted, median-filtered BV images of NGC 1850. The solid lines are isophotes while the dashed lines are the best-fit ellipses.

Fig. 11– B and V surface brightness profiles. The long-dashed lines are typical stellar profiles. The solid and short-dashed lines are isotropic and $r_a = 3r_s$ single-mass King-Michie surface density models, respectively. The solid horizontal lines indicate the background

levels.

Fig. 12– A finder chart for 51 of the 52 stars for which we have radial velocities.

Fig. 13– Radial velocity (upper panel) and velocity uncertainty (lower panel) vs. B-V.

Fig. 14– Mean radial velocity vs. projected radius (upper panel) and versus position angle (lower panel) for the 36 remaining stars. The solid lines are the mean velocity, $\bar{v} = 251.4 \pm 2.0 \text{ km s}^{-1}$.

Fig. 15– The lower left panel displays the difference in median velocity for stars on either side of an axis at the specified position angle. Also shown is the best fit sine function corresponding to a rotation axis with position angle 100° . The other panels show the same thing but for velocity data which has been rotation-subtracted assuming the specified ellipticities and the models described in §5.2.

Table 1
BV Photometry

ID	X	Y	V	B-V	ID	X	Y	V	B-V	ID	X	Y	V	B-V
1	511.8	417.2	18.04	1.04	51	505.7	444.8	18.04	-0.04	101	528.5	463.9	13.40	-0.14
2	528.8	418.0	18.75	-0.15	52	545.1	445.0	18.54	-0.14	102	498.6	464.4	14.64	0.29
3	541.4	418.1	16.37	-0.00	53	485.8	445.1	18.08	0.03	103	466.2	464.5	16.84	1.58
4	502.5	419.7	18.44	1.23	54	530.6	445.2	14.79	-0.14	104	548.6	464.5	16.06	0.11
5	521.2	419.9	16.28	-0.19	55	550.8	445.7	16.01	0.18	105	555.2	465.2	18.49	-0.03
6	545.2	420.4	18.85	-0.07	56	498.4	446.3	18.30	-0.10	106	481.9	465.3	17.01	-0.05
7	498.8	422.3	17.07	-0.02	57	492.8	446.5	18.48	0.02	107	534.1	465.3	14.01	-0.13
8	506.3	423.8	16.95	-0.02	58	516.4	446.6	16.42	0.05	108	518.2	465.9	16.98	-0.15
9	552.8	426.0	15.90	-0.05	59	463.3	447.1	19.48	-0.01	109	526.0	466.2	15.40	-0.16
10	544.8	426.5	16.82	-0.08	60	579.3	447.2	17.81	-0.03	110	607.7	466.4	17.95	-0.11
11	558.7	426.6	18.51	-0.02	61	549.8	447.9	16.87	-0.11	111	485.1	466.5	18.02	-0.00
12	533.7	426.8	18.14	0.09	62	594.6	448.0	18.78	0.00	112	542.2	466.8	15.48	-0.16
13	480.4	427.3	18.40	-0.00	63	565.7	448.0	18.81	-0.19	113	610.3	467.1	17.35	0.11
14	522.6	429.1	17.82	-0.03	64	570.4	448.9	18.53	0.38	114	480.2	467.4	17.00	0.07
15	499.4	429.4	17.30	-0.05	65	534.8	450.1	18.18	-0.08	115	477.9	467.6	17.99	-0.16
16	572.9	429.4	18.86	0.19	66	486.8	451.3	17.99	-0.07	116	562.6	468.7	16.83	-0.00
17	531.8	429.4	18.64	-0.10	67	552.1	451.5	19.08	-0.14	117	459.1	468.8	16.97	-0.00
18	565.5	430.2	18.98	-0.06	68	589.2	452.1	17.85	-0.06	118	511.2	469.1	17.17	-0.10
19	502.7	430.2	18.86	-0.20	69	463.8	452.2	17.17	-0.01	119	534.6	469.5	17.22	-0.03
20	537.0	430.3	18.84	-0.09	70	554.9	452.5	18.34	-0.11	120	584.5	469.6	17.81	-0.05
21	523.9	432.8	17.08	-0.04	71	515.3	452.6	17.91	-0.11	121	555.6	469.6	18.05	-0.14
22	579.2	434.0	18.39	-0.09	72	452.9	452.8	18.17	-0.17	122	567.2	470.3	17.54	0.02
23	569.2	435.5	16.51	-0.01	73	572.1	452.9	16.81	-0.01	123	528.1	470.5	18.41	-0.34
24	476.1	435.8	17.03	-0.01	74	494.1	453.8	16.53	1.45	124	549.2	470.6	18.45	-0.16
25	511.8	435.8	17.62	0.08	75	541.1	454.9	17.86	-0.13	125	575.1	470.8	17.11	-0.01
26	466.5	437.0	18.06	-0.05	76	491.9	455.3	15.76	1.46	126	523.4	470.9	17.77	0.12
27	504.5	437.4	18.75	0.31	77	556.4	455.6	18.94	-0.25	127	494.3	471.2	13.69	-0.14
28	490.7	437.7	18.15	0.07	78	515.5	456.9	16.28	-0.09	128	544.7	471.7	17.58	-0.10
29	471.6	437.9	15.92	-0.06	79	590.4	457.0	17.16	-0.08	129	608.3	471.7	18.34	0.07
30	539.8	437.9	16.81	0.02	80	492.2	457.1	17.09	1.47	130	439.6	471.9	17.60	0.07
31	480.3	439.0	17.24	0.95	81	542.5	457.6	15.83	-0.12	131	552.8	472.2	17.14	-0.09
32	502.0	439.2	17.62	-0.06	82	522.5	457.8	16.92	-0.08	132	445.7	472.6	18.81	-0.21
33	572.7	439.2	18.65	0.22	83	556.2	457.9	17.64	-0.05	133	530.5	473.3	18.01	-0.02
34	531.5	439.2	17.79	-0.08	84	532.6	458.0	18.18	0.18	134	473.9	473.6	16.66	-0.10
35	486.7	440.0	15.81	0.01	85	527.6	458.2	14.32	-0.16	135	513.8	473.7	15.43	1.22
36	512.0	440.3	17.78	-0.16	86	443.2	459.1	18.44	0.00	136	524.2	474.4	14.96	0.12
37	581.7	441.1	17.11	-0.04	87	521.1	459.4	15.43	-0.14	137	450.7	474.7	19.29	0.12
38	589.4	441.5	18.64	0.04	88	460.3	459.7	15.06	1.02	138	613.5	474.8	18.38	0.01
39	572.5	441.5	18.04	-0.06	89	569.7	460.4	18.61	0.03	139	459.8	475.1	17.82	0.07
40	464.3	441.7	16.90	1.46	90	487.1	460.8	17.82	0.09	140	568.6	475.6	17.63	-0.15
41	549.6	441.8	18.11	-0.04	91	539.8	461.2	19.41	-0.22	141	472.3	476.2	17.20	-0.11
42	569.3	441.8	19.04	-0.07	92	524.3	461.5	13.54	-0.13	142	550.6	476.4	16.81	0.00
43	527.8	443.0	15.39	1.28	93	492.7	461.7	17.33	-0.07	143	492.8	476.7	17.07	-0.07
44	558.1	443.1	17.75	-0.12	94	534.0	461.9	17.04	-0.06	144	457.3	477.0	17.50	1.11
45	587.1	443.6	16.20	-0.04	95	557.8	462.6	17.06	-0.06	145	466.2	477.1	17.60	-0.09
46	540.8	443.6	18.43	0.20	96	469.1	462.8	17.22	-0.03	146	494.6	477.7	17.64	-0.16
47	492.4	443.9	17.94	-0.06	97	523.6	463.2	13.09	-0.12	147	579.6	477.9	17.76	-0.02
48	523.3	444.4	18.11	0.00	98	495.5	463.2	16.72	0.72	148	532.3	478.1	18.17	-0.08
49	471.0	444.4	19.15	0.06	99	451.6	463.4	18.75	0.13	149	566.4	478.3	13.74	0.24
50	573.9	444.5	17.68	-0.05	100	579.6	463.9	17.67	0.10	150	588.6	478.5	18.13	0.63

Table 1 (cont.)
BV Photometry

ID	X	Y	V	B-V	ID	X	Y	V	B-V	ID	X	Y	V	B-V
151	559.0	478.8	17.77	-0.03	201	535.1	492.3	15.31	0.98	251	542.2	502.8	14.76	0.12
152	613.8	478.9	14.85	0.50	202	586.1	492.4	18.30	-0.06	252	469.2	503.1	17.36	-0.04
153	481.1	479.1	18.48	-0.17	203	540.5	492.7	17.08	0.10	253	488.7	503.5	18.32	0.04
154	472.8	479.4	16.73	-0.00	204	469.1	493.1	17.47	0.07	254	511.9	503.6	14.72	0.13
155	529.5	479.5	17.41	-0.12	205	572.0	493.3	18.20	0.03	255	595.2	503.8	18.15	0.14
156	562.3	480.1	18.37	-0.06	206	495.2	493.5	16.25	-0.05	256	479.4	504.0	18.11	-0.04
157	569.4	480.1	17.32	0.04	207	567.3	493.7	15.42	-0.09	257	534.4	504.3	16.86	-0.08
158	556.0	480.4	17.33	0.00	208	463.9	493.8	15.79	-0.02	258	454.3	504.4	15.65	1.24
159	485.3	480.6	18.58	-0.24	209	554.9	493.8	15.40	1.26	259	499.9	504.4	13.56	0.65
160	468.2	481.1	16.56	-0.05	210	432.3	494.2	18.53	-0.03	260	467.2	504.7	17.38	-0.05
161	591.7	481.2	18.54	-0.04	211	560.5	494.2	18.63	0.32	261	494.5	505.0	15.36	0.86
162	574.4	481.4	18.37	-0.06	212	607.0	494.2	18.31	0.01	262	484.4	505.4	15.37	1.44
163	581.5	481.8	19.10	0.03	213	437.0	494.2	14.85	0.19	263	580.5	505.6	18.60	0.31
164	587.7	482.1	16.90	0.05	214	484.6	494.3	17.68	-0.02	264	477.8	506.2	18.05	0.10
165	502.4	482.7	17.61	-0.03	215	593.2	494.8	18.78	0.05	265	447.1	506.6	16.69	-0.08
166	536.7	482.9	16.74	-0.07	216	475.9	494.8	17.99	0.03	266	552.3	507.5	18.12	-0.06
167	508.6	483.1	15.56	0.88	217	574.4	495.6	18.96	0.33	267	584.0	507.7	18.13	0.18
168	462.0	483.2	18.35	-0.12	218	523.7	495.9	16.01	0.09	268	574.3	507.9	18.12	0.04
169	442.6	483.7	18.90	-0.10	219	540.8	495.9	17.22	0.06	269	568.8	507.9	19.12	-0.13
170	598.9	484.2	13.74	-0.07	220	480.5	496.3	16.53	-0.02	270	558.2	508.4	15.81	0.10
171	435.0	484.5	19.30	-0.32	221	547.4	496.4	16.73	-0.01	271	510.1	508.6	15.86	-0.24
172	484.5	484.7	18.18	-0.03	222	508.1	496.4	15.06	0.90	272	479.1	509.0	17.12	0.03
173	539.8	484.7	14.26	0.78	223	598.0	496.7	17.95	-0.04	273	449.1	509.1	18.48	-0.03
174	545.7	485.1	17.19	0.11	224	453.2	496.7	17.95	-0.05	274	527.3	509.1	16.19	-0.17
175	532.4	485.3	15.21	1.40	225	491.2	497.0	16.62	-0.04	275	578.8	509.2	18.41	-0.02
176	478.2	485.5	17.64	0.18	226	467.5	497.2	18.17	-0.05	276	499.1	509.6	16.85	0.09
177	474.4	485.5	14.94	0.27	227	483.9	497.4	18.40	0.20	277	608.8	509.6	17.20	0.07
178	469.7	485.5	18.30	0.14	228	504.5	497.6	16.32	-0.04	278	475.2	509.7	17.50	0.00
179	613.8	485.8	18.76	0.09	229	487.2	497.7	17.42	0.93	279	554.4	509.8	17.41	-0.07
180	565.4	486.0	18.93	0.28	230	559.2	498.3	18.20	0.19	280	540.8	510.1	17.94	-0.22
181	526.8	486.0	17.16	-0.15	231	561.7	498.8	18.42	-0.03	281	491.3	510.2	15.94	-0.02
182	535.9	486.0	14.74	0.20	232	578.8	498.8	17.23	0.00	282	432.2	510.6	16.78	-0.00
183	440.6	486.3	19.15	-0.12	233	440.8	498.9	18.68	-0.02	283	604.6	511.1	14.96	0.38
184	595.7	486.4	18.07	-0.18	234	541.7	499.0	17.90	0.01	284	528.5	511.2	14.69	0.86
185	448.4	487.3	16.80	0.01	235	609.4	499.0	17.93	0.00	285	425.5	511.3	17.45	-0.02
186	463.1	487.6	17.08	0.02	236	515.0	499.0	15.86	1.01	286	587.2	511.5	18.55	0.10
187	458.5	487.8	17.94	0.03	237	572.1	499.2	18.42	0.07	287	440.1	511.6	18.18	0.03
188	561.9	487.9	18.57	0.06	238	463.8	499.6	16.29	-0.13	288	485.5	511.6	16.27	-0.07
189	468.8	488.0	18.95	-0.11	239	426.1	500.5	16.96	0.03	289	567.5	511.7	18.56	-0.16
190	504.4	488.3	16.57	-0.01	240	467.0	500.5	17.52	0.06	290	512.7	511.9	14.57	0.10
191	534.4	488.5	16.33	-0.04	241	445.7	501.3	17.96	-0.02	291	492.1	512.2	14.87	0.17
192	552.3	488.6	18.20	-0.07	242	437.6	501.4	17.78	0.10	292	599.7	512.4	16.83	0.12
193	485.0	489.7	18.23	-0.13	243	464.9	501.5	16.99	0.03	293	532.5	512.7	14.62	0.15
194	573.6	490.9	16.53	0.06	244	481.2	501.5	17.08	0.03	294	550.9	513.1	17.03	-0.28
195	482.5	491.1	17.91	-0.24	245	497.1	501.8	16.02	-0.04	295	582.4	513.1	18.50	-0.02
196	563.9	491.3	18.11	-0.01	246	430.1	502.4	15.24	0.09	296	530.1	513.3	14.85	0.18
197	613.2	491.4	18.48	0.17	247	475.6	502.5	16.25	-0.03	297	447.7	513.5	17.99	0.06
198	505.7	491.5	15.30	0.13	248	601.9	502.5	17.26	-0.03	298	591.5	513.5	18.53	0.09
199	448.4	491.6	18.77	0.25	249	535.8	502.6	16.70	-0.12	299	573.2	513.5	18.63	-0.15
200	530.6	492.0	17.28	0.01	250	574.4	502.7	15.47	1.71	300	615.0	513.6	17.80	-0.03

Table 1 (cont.)
BV Photometry

ID	X	Y	V	B-V	ID	X	Y	V	B-V	ID	X	Y	V	B-V
301	542.6	513.9	16.48	0.01	351	560.8	526.2	15.39	1.09	401	499.4	538.7	16.38	0.03
302	488.2	513.9	14.76	1.21	352	538.0	526.3	15.40	-0.00	402	535.9	538.7	15.33	0.01
303	426.7	514.4	18.95	-0.20	353	575.7	526.4	18.32	0.07	403	547.4	539.0	17.33	-0.16
304	480.5	514.5	15.46	1.53	354	471.4	526.5	19.00	0.07	404	557.1	539.2	17.01	-0.10
305	423.9	514.8	17.40	0.04	355	540.7	526.7	16.60	-0.07	405	528.4	539.4	17.88	0.22
306	434.4	515.2	18.19	0.23	356	520.2	526.9	14.43	0.31	406	552.5	539.5	16.86	0.01
307	602.5	515.3	17.80	0.16	357	509.0	527.4	17.81	0.45	407	571.8	539.7	18.47	0.08
308	451.4	515.4	17.97	0.16	358	552.0	528.2	18.13	0.29	408	602.8	540.0	18.01	0.20
309	559.7	515.5	16.57	0.16	359	466.5	528.2	19.05	-0.08	409	443.4	540.0	18.02	0.06
310	583.6	515.6	17.95	0.00	360	580.5	528.2	18.81	0.10	410	437.4	540.7	18.71	-0.04
311	576.4	515.6	18.07	0.09	361	555.0	528.3	17.76	0.17	411	524.8	540.7	16.93	0.01
312	462.4	515.9	18.75	-0.17	362	613.1	528.6	15.44	2.08	412	612.0	540.9	19.13	0.62
313	550.7	516.2	15.70	-0.11	363	480.7	528.7	17.84	-0.09	413	483.6	541.0	18.59	-0.17
314	540.9	516.6	17.23	0.35	364	517.3	528.8	16.63	0.03	414	435.2	541.2	17.97	0.05
315	497.6	517.0	17.58	-0.04	365	558.2	529.1	16.83	0.12	415	443.5	542.2	18.91	0.01
316	558.6	517.0	17.19	0.07	366	473.2	529.1	16.95	0.04	416	495.7	542.2	17.67	0.04
317	580.7	517.0	16.35	1.29	367	535.2	529.3	14.53	0.17	417	588.5	542.5	17.24	-0.31
318	530.2	517.7	15.44	0.63	368	566.6	530.0	17.73	-0.09	418	530.9	543.0	16.70	0.06
319	542.5	518.4	16.77	0.06	369	501.5	530.5	15.68	1.64	419	511.7	543.3	17.15	-0.12
320	477.0	518.5	14.97	0.19	370	569.7	531.2	15.64	1.46	420	556.2	543.5	17.24	0.17
321	494.9	518.5	17.78	0.09	371	467.4	531.2	18.48	0.29	421	609.8	544.0	19.23	0.12
322	488.0	519.2	15.34	1.24	372	589.8	531.5	18.32	0.02	422	589.9	544.1	14.84	1.75
323	566.0	519.2	17.21	0.13	373	498.7	532.1	17.43	-0.25	423	571.7	544.4	18.28	0.14
324	501.5	519.3	15.58	0.30	374	561.8	532.2	16.87	0.02	424	558.7	544.9	15.36	1.55
325	555.5	519.4	17.29	-0.24	375	508.3	532.3	17.35	0.06	425	564.4	545.3	18.16	0.20
326	481.0	519.4	18.11	-0.03	376	539.9	532.5	16.48	0.34	426	515.5	546.6	15.60	1.34
327	551.5	519.7	17.53	-0.49	377	514.5	533.0	15.56	1.31	427	569.2	547.0	16.03	0.10
328	575.3	519.8	17.61	0.09	378	528.9	533.2	15.39	1.42	428	441.4	547.3	17.96	0.02
329	457.3	520.2	17.62	0.10	379	517.2	533.3	15.78	0.03	429	520.1	547.7	17.21	0.00
330	610.8	520.4	14.99	0.31	380	556.8	533.5	18.08	0.03	430	510.9	548.0	17.16	0.02
331	543.1	520.6	17.60	-0.05	381	560.0	533.6	16.84	-0.05	431	559.6	548.0	16.94	-0.01
332	484.4	520.7	15.86	1.33	382	548.6	534.0	14.81	0.19	432	471.2	548.1	17.92	0.04
333	498.2	520.7	15.86	1.24	383	526.3	534.0	16.70	-0.16	433	488.0	548.2	14.94	1.22
334	433.3	520.9	18.10	-0.15	384	539.2	534.6	15.27	0.15	434	434.0	548.5	18.32	-0.11
335	505.2	521.6	16.74	0.24	385	572.9	535.0	15.79	0.06	435	475.9	548.5	18.32	-0.26
336	467.7	521.8	15.56	1.22	386	466.8	535.0	18.52	-0.03	436	499.9	548.8	17.72	-0.10
337	443.7	522.2	15.13	1.70	387	595.0	535.5	16.90	0.01	437	528.8	549.9	17.36	0.07
338	478.7	522.5	15.70	1.58	388	483.2	535.6	15.64	1.27	438	586.5	550.2	18.33	0.21
339	564.5	522.9	17.22	-0.02	389	504.1	535.7	18.06	0.03	439	547.5	550.2	16.45	0.03
340	585.2	522.9	17.55	0.08	390	617.9	536.1	19.36	0.21	440	451.8	550.7	18.12	0.01
341	556.8	522.9	14.89	0.24	391	570.5	536.2	15.41	1.43	441	559.5	550.7	15.71	1.28
342	447.6	523.0	18.07	0.11	392	510.1	536.2	17.38	0.05	442	551.0	550.7	17.89	0.15
343	536.9	523.1	17.87	-0.12	393	543.0	536.4	14.55	0.25	443	515.0	550.9	17.16	-0.09
344	509.2	523.2	16.30	0.04	394	560.2	536.5	14.91	0.25	444	491.5	551.1	18.06	0.08
345	611.0	524.2	17.34	0.42	395	491.0	536.7	18.66	0.24	445	540.2	551.2	17.22	-0.72
346	559.6	524.3	14.53	0.11	396	556.7	536.9	14.83	0.19	446	533.4	551.2	17.36	0.18
347	486.8	524.5	17.24	0.00	397	435.0	537.1	18.29	0.08	447	432.1	551.3	18.52	0.15
348	516.6	524.6	15.97	0.13	398	427.1	538.0	17.85	-0.01	448	553.5	551.6	17.01	0.02
349	505.5	525.5	15.59	1.05	399	600.6	538.3	18.00	0.24	449	455.8	551.6	17.14	0.00
350	442.8	526.2	18.51	0.04	400	524.0	538.3	14.99	0.05	450	565.8	552.5	17.79	0.02

Table 1 (cont.)
BV Photometry

ID	X	Y	V	B-V	ID	X	Y	V	B-V	ID	X	Y	V	B-V
451	571.9	552.6	18.65	0.00	501	497.5	564.2	13.68	0.23	551	483.9	578.0	18.48	-0.05
452	470.0	552.7	18.05	0.26	502	484.0	565.6	17.61	-0.02	552	505.1	578.9	17.34	-0.03
453	534.5	553.1	17.35	-0.13	503	471.4	566.3	18.08	0.02	553	494.5	579.1	16.59	-0.00
454	544.1	553.2	17.38	-0.14	504	527.4	566.6	17.36	0.02	554	564.9	579.4	19.23	0.03
455	578.6	553.6	17.81	0.16	505	607.7	566.6	16.79	1.65	555	554.7	579.4	16.89	-0.01
456	463.0	553.7	18.96	-0.21	506	562.8	566.6	18.17	-0.01	556	577.4	579.7	18.72	0.29
457	481.0	554.2	16.42	-0.12	507	595.6	567.3	18.46	0.22	557	487.6	580.2	18.59	0.11
458	527.8	554.2	15.96	1.34	508	574.6	567.4	16.94	0.09	558	509.2	580.2	18.63	0.07
459	457.2	554.3	16.76	-0.05	509	585.8	567.6	16.06	1.48	559	519.5	580.8	16.89	0.08
460	531.7	554.5	17.02	-0.10	510	441.2	567.8	14.87	0.11	560	469.2	581.8	15.32	0.14
461	545.5	554.8	17.61	-0.17	511	556.0	567.9	18.87	0.41	561	556.2	582.1	18.82	-0.05
462	559.3	555.2	17.63	0.15	512	486.7	568.2	18.29	0.01	562	575.5	582.2	17.63	0.03
463	460.4	555.6	17.18	0.08	513	483.2	569.0	18.13	0.06	563	462.8	582.2	18.28	-0.02
464	485.8	555.6	17.69	0.04	514	520.5	569.1	14.68	0.30	564	581.4	582.3	18.50	0.12
465	448.6	555.7	18.83	0.05	515	462.7	569.3	19.03	-0.25	565	505.0	583.3	18.81	0.12
466	435.8	555.8	19.18	0.13	516	470.6	569.9	18.41	0.07	566	511.0	584.0	15.60	1.02
467	529.0	555.9	15.90	1.33	517	465.2	570.0	17.34	0.18	567	473.9	584.6	17.88	0.07
468	611.1	556.1	19.25	0.09	518	541.8	570.0	17.73	0.10	568	561.5	585.4	17.70	0.84
469	509.6	556.4	17.78	-0.00	519	561.4	570.3	18.12	-0.06	569	460.4	585.7	17.28	0.16
470	478.5	556.5	15.40	0.00	520	546.8	570.3	17.76	-0.04	570	523.5	585.8	19.06	-0.20
471	506.1	556.7	15.46	1.63	521	479.0	570.3	17.82	-0.26	571	514.3	585.9	18.76	0.30
472	493.9	556.7	17.72	-0.08	522	598.3	570.3	18.06	0.05	572	493.4	586.0	19.07	-0.34
473	482.3	557.2	18.22	0.07	523	514.1	570.5	18.15	0.13	573	549.5	586.3	18.42	0.14
474	569.7	557.5	17.92	0.11	524	584.9	571.2	17.95	0.13	574	496.6	586.8	17.85	0.12
475	600.8	557.7	19.09	0.13	525	553.0	571.4	18.88	0.10	575	481.8	587.1	18.55	0.08
476	540.2	557.7	17.84	-0.08	526	481.9	571.5	18.59	0.01	576	457.7	587.2	16.07	1.41
477	498.1	557.8	17.77	-0.08	527	451.8	572.0	19.15	0.05	577	486.0	587.4	17.25	0.23
478	543.0	558.5	17.82	0.09	528	496.8	572.2	16.53	0.01	578	500.5	587.5	18.73	0.14
479	551.6	558.8	17.92	0.05	529	475.6	572.2	16.88	-0.34	579	544.1	587.8	18.06	-0.08
480	486.8	559.2	16.39	1.43	530	557.1	572.2	18.15	-0.07	580	588.2	588.1	19.18	0.04
481	526.7	559.8	18.47	0.06	531	504.1	572.5	18.76	-0.22	581	490.8	588.1	16.25	0.42
482	582.0	560.1	17.55	0.06	532	471.6	572.6	16.54	0.08	582	502.7	588.9	16.98	0.51
483	474.2	560.2	19.02	0.08	533	499.3	572.9	16.15	-0.08	583	549.2	589.2	18.07	0.01
484	523.2	560.6	18.32	-0.03	534	600.2	573.0	18.59	0.30	584	505.7	589.8	17.76	-0.01
485	485.2	560.6	16.07	0.64	535	468.0	573.6	18.01	0.24	585	529.0	589.9	18.84	0.12
486	500.5	560.9	17.94	0.03	536	528.7	573.9	18.52	-0.13	586	478.2	590.0	18.23	0.02
487	536.0	561.0	16.70	-0.04	537	458.4	574.0	18.87	0.18	587	583.6	590.0	15.73	1.54
488	446.7	562.2	18.14	0.09	538	548.6	574.1	19.24	-0.10	588	522.7	590.6	18.99	0.01
489	559.7	562.3	14.58	1.70	539	486.9	574.4	19.00	0.15	589	514.0	590.9	19.05	0.06
490	512.3	562.6	18.16	0.11	540	521.7	574.5	18.58	-0.53	590	539.9	591.2	17.50	-0.07
491	588.0	562.7	17.81	0.02	541	535.9	574.6	18.29	-0.13	591	576.7	591.3	19.01	-0.09
492	533.0	562.8	17.98	0.08	542	493.2	574.9	19.36	0.10	592	489.2	591.3	15.50	1.11
493	480.6	563.1	18.37	-0.03	543	454.5	575.4	18.41	0.02	593	509.4	591.5	18.40	0.08
494	502.5	563.2	18.03	0.08	544	477.4	575.5	19.53	-0.15	594	553.8	592.3	17.77	-0.04
495	443.8	563.6	17.23	-0.07	545	573.3	575.5	17.72	0.02	595	517.3	592.7	18.32	-0.01
496	542.5	563.8	16.15	1.48	546	553.6	575.6	16.27	0.92	596	504.4	592.7	18.14	-0.01
497	551.4	564.0	15.18	0.39	547	543.0	576.6	18.14	-0.03	597	570.4	593.1	14.85	0.17
498	530.6	564.0	16.07	0.02	548	497.2	577.2	18.31	0.21	598	477.3	593.5	16.78	0.07
499	606.1	564.0	17.81	0.31	549	546.6	577.6	17.27	0.03	599	486.6	593.8	16.75	1.70
500	508.9	564.2	17.17	-0.02	550	463.3	577.9	18.75	0.56	600	583.1	594.1	15.29	1.14

Table 1 (cont.)
BV Photometry

ID	X	Y	V	B-V	ID	X	Y	V	B-V	ID	X	Y	V	B-V
601	473.9	594.3	17.55	0.09	616	566.0	598.7	19.08	-0.36	631	518.5	604.1	19.30	0.49
602	471.3	594.5	17.59	0.01	617	479.5	599.1	18.99	0.24	632	509.2	604.3	16.61	0.05
603	509.7	594.7	17.49	0.12	618	539.2	599.1	18.05	0.12	633	525.5	604.9	18.08	0.03
604	498.9	594.8	17.65	0.28	619	529.0	599.4	18.00	0.17	634	484.8	605.1	16.65	-0.04
605	555.9	595.1	17.52	-0.03	620	542.3	599.6	18.43	0.11	635	499.6	605.4	14.69	0.39
606	546.7	595.4	18.71	0.04	621	559.2	599.6	15.51	0.28	636	492.9	606.0	17.60	0.05
607	519.1	595.5	18.01	-0.05	622	473.7	599.7	17.63	0.01	637	540.7	606.0	18.26	-0.04
608	552.3	595.7	17.67	0.12	623	514.1	600.1	18.34	0.09	638	553.8	606.3	17.18	0.01
609	532.4	596.1	18.60	0.03	624	485.8	601.7	17.69	1.15	639	485.3	607.3	17.06	0.12
610	508.5	596.3	17.93	0.09	625	496.6	602.0	16.64	0.14	640	512.4	607.7	18.30	-0.06
611	564.2	597.0	18.48	-0.04	626	477.4	602.3	16.82	0.66	641	504.8	610.1	17.95	0.10
612	492.4	597.1	15.86	1.27	627	475.6	602.5	17.15	0.10	642	528.8	612.0	18.30	1.04
613	483.0	597.1	18.22	0.05	628	491.8	602.7	18.27	-0.14	643	534.9	614.7	18.06	-0.10
614	486.4	597.9	18.39	0.54	629	482.0	603.0	18.59	0.16					
615	500.9	598.5	19.30	-0.01	630	494.3	603.9	17.67	0.02					

Table 2
Surface Photometry

R (pc)	L_B ($L_{B\odot} \text{ pc}^{-2}$)	R (pc)	L_V ($L_{V\odot} \text{ pc}^{-2}$)
0.3	26246.0 ± 3517.0	0.3	14938.0 ± 1958.0
0.5	25094.0 ± 3993.0	0.5	9719.0 ± 1844.0
0.7	24901.0 ± 3821.0	0.7	10536.0 ± 2155.0
0.8	31223.0 ± 5737.0	0.8	12438.0 ± 3273.0
1.1	26021.0 ± 6693.0	1.0	11899.0 ± 3994.0
1.3	22898.0 ± 4908.0	1.3	11771.0 ± 2702.0
1.7	19512.0 ± 4085.0	1.7	10236.0 ± 2041.0
2.1	14890.0 ± 3159.0	2.1	6037.0 ± 1010.0
2.6	11276.0 ± 1308.0	2.7	5831.0 ± 988.0
3.3	10828.0 ± 1290.0	3.3	5357.0 ± 554.0
4.2	6910.0 ± 1417.0	4.2	3517.0 ± 733.0
5.3	3889.0 ± 938.0	5.3	2098.0 ± 443.0
6.6	3198.0 ± 435.0	6.6	1625.0 ± 261.0
8.4	2919.0 ± 429.0	8.4	1423.0 ± 199.0
10.5	1044.0 ± 210.0	10.5	510.0 ± 88.0
13.2	944.0 ± 201.0	13.2	423.0 ± 101.0
16.7	414.0 ± 144.0	16.7	233.0 ± 68.0
21.0	360.0 ± 63.0	21.0	170.0 ± 29.0
26.4	140.0 ± 44.0	26.4	64.0 ± 23.0
33.1	92.0 ± 39.0	33.1	46.0 ± 20.0
40.5	22.0 ± 28.0	40.5	8.0 ± 17.0

Table 3
King-Michie - B Band Fitted Parameters

	Photometry					Velocities		
r_a (r_s)	W_o	r_s (pc)	c	χ^2_ν ($\nu = 18$)	$P(> \chi^2_\nu)$	v_s (km s $^{-1}$)	ζ^2	$P(> \zeta^2)$
ISO	7.3 ± 0.30	2.6 ± 0.20	$42. \pm 8.$	0.68	0.81	2.2 ± 0.4	37.61	0.21
20	7.3 ± 0.30	2.6 ± 0.20	$46. \quad ^{+12.}_{-9.}$	0.69	0.80	2.2 ± 0.4	37.66	0.21
10	7.3 ± 0.35	2.6 ± 0.15	$67. \quad ^{+45.}_{-25.}$	0.72	0.76	2.2 ± 0.4	37.80	0.20
5	$6.8 \quad ^{+0.10}_{-0.20}$	3.1 ± 0.15	$125. \pm 50.$	0.86	0.60	2.2 ± 0.5	38.40	0.15
3	$5.9 \quad ^{+0.05}_{-0.10}$	3.5 ± 0.20	$122. \pm 45.$	1.17	0.25	2.3 ± 0.5	39.11	0.10

Table Table 4
King-Michie - V Band Fitted Parameters

	Photometry					Velocities		
r_a (r_s)	W_o	r_s (pc)	c	χ^2_ν ($\nu = 18$)	$P(> \chi^2_\nu)$	v_s (km s $^{-1}$)	ζ^2	$P(> \zeta^2)$
ISO	7.1 ± 0.30	2.7 ± 0.20	$37. \pm 8.$	0.91	0.51	2.2 ± 0.4	37.68	0.21
20	7.1 ± 0.30	2.8 ± 0.20	$39. \quad ^{+11.}_{-9.}$	0.91	0.51	2.2 ± 0.4	37.72	0.20
10	7.1 ± 0.35	2.8 ± 0.20	$51. \quad ^{+33.}_{-16.}$	0.92	0.50	2.2 ± 0.4	37.84	0.19
5	$6.8 \quad ^{+0.10}_{-0.20}$	3.2 ± 0.20	$116. \pm 47.$	0.98	0.43	2.2 ± 0.5	38.31	0.15
3	$5.9 \quad ^{+0.05}_{-0.10}$	3.7 ± 0.20	$122. \pm 44.$	1.16	0.25	2.3 ± 0.5	38.97	0.11

Table 5
King-Michie - B Band Derived Parameters

r_a (r_s)	$\rho_{KB\odot}$ ($L_{B\odot} \text{ pc}^{-3}$)	L_B ($10^6 L_{B\odot}$)	Mass ($10^4 M_\odot$)	M/L_B ($M_\odot/L_{B\odot}$)
ISO	$5500. \pm 700.$	2.60 ± 0.20	5.7 ± 2.3	0.02 ± 0.01
20	$5500. \pm 700.$	2.64 ± 0.20	5.7 ± 2.3	0.02 ± 0.01
10	$5400. \pm 700.$	2.75 ± 0.30	5.9 ± 2.4	0.02 ± 0.01
5	$4200. \pm 550.$	2.75 ± 0.20	5.8 ± 2.3	0.02 ± 0.01
3	$3700. \pm 500.$	2.62 ± 0.20	5.4 ± 2.2	0.02 ± 0.01

Table 6
King-Michie - V Band Derived Parameters

r_a (r_s)	$\rho_{KV\odot}$ ($L_{V\odot} pc^{-3}$)	L_V ($10^6 L_{V\odot}$)	Mass ($10^4 M_\odot$)	M/ L_V ($M_\odot/L_{V\odot}$)
ISO	2300. $\pm$ 350.	1.26 $\pm$ 0.10	5.7 $\pm$ 2.3	0.05 $\pm$ 0.02
20	2300. $\pm$ 350.	1.27 $\pm$ 0.10	5.7 $\pm$ 2.3	0.04 $\pm$ 0.02
10	2300. $\pm$ 300.	1.31 $\pm$ 0.15	5.8 $\pm$ 2.3	0.04 $\pm$ 0.02
5	1900. $\pm$ 250.	1.37 $\pm$ 0.10	5.9 $\pm$ 2.4	0.04 $\pm$ 0.02
3	1600. $\pm$ 200.	1.32 $\pm$ 0.10	5.6 $\pm$ 2.2	0.04 $\pm$ 0.02

Table 7
Velocity Standards

HJD (-2448000)	v_x (km s ⁻¹)	$\langle v_x \rangle$ (km s ⁻¹)	χ^2_ν
<u>Local</u>			
304.5946	34.5 ± 0.3	34.2 ± 0.1	1.12
305.5481	34.3 ± 0.3		
306.5467	33.7 ± 0.3		
307.5481	34.4 ± 0.2		
308.5432	34.0 ± 0.3		
605.7090	36.0 ± 0.6	35.7 ± 0.3	0.72
606.6605	36.0 ± 0.6		
607.6073	35.0 ± 0.6		
608.6622	35.7 ± 0.6		
<u>HD 23214</u>			
305.5204	-2.6 ± 0.4	-2.9 ± 0.2	0.26
306.5211	-2.9 ± 0.4		
307.5189	-3.0 ± 0.4		
308.5182	-3.1 ± 0.4		
605.5891	-3.1 ± 0.5	-2.8 ± 0.2	0.32
605.6899	-2.8 ± 0.6		
606.5868	-2.9 ± 0.6		
607.5920	-2.4 ± 0.5		
608.5895	-2.7 ± 0.5		

Table 8
Radial Velocities

ID	R (pc)	Θ ($^{\circ}$)	v_x (km s $^{-1}$)	$\langle v_x \rangle$ (km s $^{-1}$)	HJD (-2448000)	V (mag)	B-V (mag)
RV1	0.0	18.5	252.7 ± 2.1	252.7 ± 2.1	608.6797	15.74	0.58
RV2	0.5	166.7	248.3 ± 1.3	248.3 ± 1.3	608.6768	14.89	0.20
RV3	0.9	224.6	248.4 ± 1.9	248.4 ± 1.9	304.5439	14.69	0.86
RV4	1.0	163.4	251.8 ± 2.6	251.8 ± 2.6	608.6729	15.44	0.67
RV5	1.1	325.4	248.4 ± 4.9	248.4 ± 4.9	303.5889	14.04	0.06
RV6	1.3	200.3	249.8 ± 3.3	249.8 ± 3.3	303.6309	14.68	0.14
RV7	1.5	71.1	252.2 ± 1.9	252.2 ± 1.9	303.5547	14.30	-0.19
RV8	1.7	349.2	260.2 ± 4.3	260.2 ± 4.3	304.5420	14.54	0.14
RV9	1.7	102.0	254.3 ± 1.7	252.3 ± 1.1	607.6504	14.23	0.68
			251.0 ± 1.4		608.6973		
RV10	2.4	321.1	257.4 ± 3.0	257.4 ± 3.0	306.5625	14.72	0.13
RV11	2.7	134.0	276.0 ± 2.8	276.0 ± 2.8	303.5850	14.53	0.18
RV12	2.8	110.9	250.3 ± 1.1	250.1 ± 0.8	607.6543	15.39	1.40
			249.9 ± 1.2		608.7061		
RV13	2.8	340.9	254.6 ± 1.3	254.6 ± 1.3	608.6816	15.14	0.60
RV14	3.0	67.9	253.1 ± 2.2	253.1 ± 2.2	608.6943	15.56	1.30
RV15	3.0	236.9	246.7 ± 1.6	246.7 ± 1.6	607.6572	15.37	0.04
RV16	3.1	76.5	248.5 ± 2.2	248.5 ± 2.2	304.5586	14.74	0.42
RV17	3.3	218.8	243.4 ± 3.1	243.4 ± 3.1	304.5479	14.73	0.13
RV18	3.6	315.3	234.4 ± 2.4	234.4 ± 2.4	306.5605	15.05	0.93
RV19	3.8	341.1	272.7 ± 1.3	260.7 ± 1.0	303.5586	13.54	0.65
RV19			240.9 ± 1.7		607.6416		
RV20	4.0	39.9	253.6 ± 1.3	253.2 ± 0.9	608.6885	15.70	1.56
			252.9 ± 1.2		605.7393		
RV21	4.1	84.4	256.7 ± 1.4	256.7 ± 1.4	605.7344	15.19	1.62
RV22	4.2	137.2	247.7 ± 1.7	247.7 ± 1.7	306.5576	14.55	0.23
RV23	4.6	147.9	258.7 ± 3.4	258.7 ± 3.4	303.5752	14.82	0.16
RV24	4.6	252.5	253.0 ± 2.0	250.1 ± 1.0	304.5547	14.71	0.21
			249.1 ± 1.2		608.6689		
RV25	4.7	0.1	242.4 ± 2.9	242.4 ± 2.9	307.5625	14.85	0.15

Table 8 (cont.)
Radial Velocities

ID	R (pc)	Θ ($^{\circ}$)	V_x (km s $^{-1}$)	$\langle v_x \rangle$ (km s $^{-1}$)	HJD (-2448000)	V (mag)	B-V (mag)
RV26	5.0	247.4	242.3 ± 2.9	242.3 ± 2.9	304.5518	14.25	0.77
RV27	5.1	57.9	248.3 ± 1.6	248.3 ± 1.6	608.6924	15.57	0.41
RV28	5.2	300.6	255.7 ± 2.1	255.7 ± 2.1	608.7168	15.55	0.86
RV29	5.2	3.4	250.5 ± 1.0	249.8 ± 0.7	608.6855	14.77	1.18
			249.2 ± 0.9		307.5654		
RV30	5.5	219.5	253.5 ± 1.5	253.5 ± 1.5	608.7139	15.40	1.23
RV31	6.0	351.4	252.2 ± 1.7	252.2 ± 1.7	606.6895	15.39	1.41
RV32	6.3	288.7	250.7 ± 2.0	250.7 ± 2.0	606.6836	15.42	1.19
RV33	6.7	14.6	251.9 ± 1.3	251.9 ± 1.3	605.7305	15.70	1.53
RV34	6.9	9.4	251.9 ± 2.0	251.9 ± 2.0	605.7266	14.96	0.19
RV35	7.2	48.0	249.7 ± 1.5	249.7 ± 1.5	607.6465	14.94	1.19
RV36	7.7	198.9	250.9 ± 1.1	250.9 ± 1.1	605.7227	15.48	1.64
RV37	7.8	309.1	277.3 ± 3.9	277.3 ± 3.9	303.5625	13.67	-0.14
RV38	8.0	91.7	241.7 ± 1.6	241.7 ± 1.6	303.6201	14.67	0.29
RV39	8.2	67.0	233.5 ± 1.1	233.2 ± 1.0	303.6152	13.67	0.23
			232.2 ± 2.0		607.6436		
RV40	8.3	226.0	239.6 ± 3.8	239.6 ± 3.8	303.5508	13.73	0.22
RV41	8.3	12.0	252.5 ± 1.3	252.5 ± 1.3	608.7246	15.57	1.25
RV42	8.3	301.6	252.5 ± 2.0	252.5 ± 2.0	308.5596	14.64	0.29
RV43	8.7	132.5	249.1 ± 1.1	249.1 ± 1.1	607.6270	14.58	1.67
RV44	10.3	356.5	250.1 ± 1.7	250.1 ± 1.7	608.7207	15.65	1.21
RV45	10.6	161.5	250.7 ± 1.0	250.7 ± 1.0	605.7178	14.85	1.73
RV46	11.8	10.2	252.4 ± 1.5	252.4 ± 1.5	606.6943	15.13	1.66
RV47	13.7	80.2	249.6 ± 1.5	249.6 ± 1.5	305.5654	14.68	0.39
RV48	16.6	304.7	252.0 ± 1.2	252.0 ± 1.2	607.6318	15.19	1.55
RV49	18.3	120.6	288.6 ± 1.1	286.3 ± 0.7	607.6240	13.66	1.82
			284.4 ± 1.0		308.5518		
RV50	18.7	67.2	253.3 ± 1.9	253.3 ± 1.9	305.5586	14.54	0.49
RV51	19.4	332.6	246.4 ± 2.6	246.4 ± 2.6	607.6377	15.08	1.41
RV52	35.5	81.7	254.9 ± 1.0	254.9 ± 1.0	307.5576	14.59	1.71

Table 9
Velocity Residuals

Iteration	v_s (km s ⁻¹)	δ_{max}	P(> δ_{max})	Star
1	4.8	4.35	0.00	39
2	2.8	3.39	0.01	38
3	2.1	2.49	0.39	52

Table 10
Rotating Ellipsoidal Models

ϵ	Mass (10 ⁴ M _⊙)	M/L _B (M _⊙ /L _{B⊙})	M/L _V (M _⊙ /L _{V⊙})	ζ^2	P(> ζ^2)	A (km s ⁻¹)
0.1	5.5 ± 0.2	0.02 ± 0.01	0.04 ± 0.02	37.47	0.34	0.9
0.2	5.6 ± 0.2	0.02 ± 0.01	0.04 ± 0.02	38.76	0.37	0.5
0.3	5.5 ± 0.2	0.02 ± 0.01	0.04 ± 0.02	40.58	0.20	0.3

Table 11
Mass Functions

M/L =		0.02 $\pm$ 0.01 M $_{\odot}$ /L $_{B\odot}$			0.05 $\pm$ 0.02 M $_{\odot}$ /L $_{V\odot}$		
m_l	m_d	x	$< m >$		x	$< m >$	
(M $_{\odot}$)	(M $_{\odot}$)		(M $_{\odot}$)			(M $_{\odot}$)	
0.05	0.0	0.29 $^{+0.3}_{-0.8}$	0.75		0.71 $^{+0.2}_{-0.4}$	0.37	
0.05	0.3	0.31 $^{+0.3}_{-0.8}$	0.83		0.79 $^{+0.2}_{-0.4}$	0.50	
0.05	0.5	0.32 $^{+0.3}_{-0.9}$	0.87		0.85 $^{+0.3}_{-0.5}$	0.57	
0.05	1.0	0.38 $^{+0.5}_{-1.0}$	0.93		1.04 $^{+0.4}_{-0.6}$	0.69	
0.05	1.5	0.46 $^{+0.5}_{-1.2}$	0.97		1.28 $^{+0.5}_{-0.8}$	0.77	
0.1	0.0	0.32 $^{+0.3}_{-0.8}$	0.93		0.77 $^{+0.2}_{-0.4}$	0.54	
0.1	0.3	0.33 $^{+0.3}_{-0.8}$	0.98		0.81 $^{+0.2}_{-0.4}$	0.62	
0.1	0.5	0.35 $^{+0.4}_{-0.9}$	1.01		0.88 $^{+0.3}_{-0.5}$	0.69	
0.1	1.0	0.41 $^{+0.5}_{-1.0}$	1.08		1.06 $^{+0.4}_{-0.6}$	0.82	
0.1	1.5	0.48 $^{+0.6}_{-1.2}$	1.13		1.30 $^{+0.5}_{-0.8}$	0.90	
0.15	0.0	0.34 $^{+0.3}_{-0.8}$	1.06		0.81 $^{+0.2}_{-0.4}$	0.66	
0.15	0.3	0.35 $^{+0.3}_{-0.8}$	1.08		0.84 $^{+0.2}_{-0.5}$	0.71	
0.15	0.5	0.37 $^{+0.4}_{-0.9}$	1.12		0.90 $^{+0.3}_{-0.5}$	0.78	
0.15	1.0	0.43 $^{+0.4}_{-1.0}$	1.19		1.08 $^{+0.4}_{-0.6}$	0.91	
0.15	1.5	0.51 $^{+0.6}_{-0.2}$	1.23		1.32 $^{+0.5}_{-0.8}$	1.00	